

Arlington Red Clover



Arlington was developed through joint efforts of the USDA Agricultural Experiment Station and Research Division of the University of Wisconsin-Madison.

Arlington is resistant to bean and yellow mosaic virus and is moderately resistant to northern anthracnose and powdery mildew.

It has a good winter survival and persists in the field longer than other varieties.



**Seeding Rate:
12-14 lbs. per acre/alone
6-8 lbs per acre/mixture**



2541 Commerce St
La Crosse WI 54603

(608)783-9560
(608)783-9515 fax
(800)328-1909 watts
www.lftseed.com

TECHNICAL DESCRIPTION



Red Clover (*Trifolium pretense* L.)

Arlington

Arlington was developed through joint efforts of the USDA Agricultural Experiment Station and Research Division of the University of Wisconsin-Madison. Arlington is resistant to bean and yellow mosaic virus and is moderately resistant to northern anthracnose and powdery mildew. It has good winter survival and persists in the field longer than other varieties.

Agronomic Traits:

Early Seedling Vigor	Excellent
Growth Habit:	Upright from Crown
Recovery After Cutting:	Very Fast
Type of Stem:	Hollow, hairy stems & branches
Stem Length:	18 inch avg.
Branches per stem:	4
Leaf Description:	Slender stalk with 3 leaflets
Taproot:	Extensively branched
Flowers:	Compact Cluster
Flower Color:	rose-pink
Seed Pods:	small & short
Seed Shape:	Kidney
Seed color:	yellow to deep violet

Crop Use Information:

Life Cycle:	Perennial
Ease of Establishment:	Fair-Good
Shade Tolerance:	Good
Drought Stress:	Excellent
Wet Soil:	Fair-Good
Low pH Tolerance:	Poor
Minimum pH:	6.0

Planting Rates:

Bushel Weight:	60 lb
Seeds Per Pound: (Non-Coated)	272,000

Rate (Lbs):	<u>Pure</u>	<u>Coated</u>	<u>With Grass</u>
North:	12-14	12-14	6-8

Seeds/Sq Ft (Non-coated)	75-88	75-88	38-50
-----------------------------	-------	-------	-------

Quality Data—Arlington Red Clover:

Adaptation and Distribution

Red clover grows best on well-drained loamy soils, but it will also grow on soil that is not as well-drained. Medium and fine textured soils are preferred by the plant over sandy or gravelly soils. It is best adapted to a pH of 6.0 or higher. Red clover is distributed throughout the United States and Canada.

Establishment

Red clover may be seeded in pure stands, but it is often mixed with grain or grass. Spring or late summer seedings are satisfactory. It may be overseeded in the spring or fall. Red clover seed should be inoculated. Phosphorus and potash are the fertilizer elements needed mostly by red clover. Apply as recommended by soil tests. Seeding may be done with a drill or broadcast. A firm, weed-free seedbed is essential. Plant seeds $\frac{1}{4}$ to $\frac{1}{2}$ inch deep. Seeding rates are 12 to 15 lbs. per acre broadcast and 6 to 8 lbs. per acre when drilled. For renovating pastures, the recommended seeding rate is 8 lbs. per acre.

Management

Graze or cut for hay when the red clover is $\frac{1}{4}$ to $\frac{1}{2}$ in bloom. A second cutting or successive grazing should occur when red clover is $\frac{1}{4}$ in bloom. Leave at least 2 to 3 inches of growth after each harvest. Care should be taken to eliminate or appreciably reduce bloating of livestock when grazing. Keep lime and fertilizers (phosphorus and potash) at the proper level.